

## THE MOST EFFECTIVE WASP & HORNET SPRAY AVAILABLE

- Drops wasps, hornets and yellow jackets instantly
- Kills with residual action as wasps and hornets return to their nests
- Jet spray up to 20 feet
- EPA Registered
- Dielectric breakdown voltage of 47,300 volts

### Description

Nu-Kill II doesn't just make wasps mad... it kills them! It is formulated with both tetramethrin and permethrin for instant kill and it provides residual kill as insects return to the nest. It does not contain any CFC's and it has a very high dielectric strength.

### Application

It instantly and effectively kills wasps, hornets, and yellow jackets. It is also excellent against crawling insects such as ants, clover mites, crickets, earwigs, millipedes, and sowbugs (pillbugs), cockroaches (including Asian cockroaches), crickets, centipedes, silverfish, ants, lone star ticks, dog ticks, crickets, fleas, earwigs, and more!

### Packaging


13.5 ounce can                      **4292-75**

### Directions for Use

**WASPS, HORNETS, YELLOW JACKETS:** For best results, spray in early morning or late evening when insect activity is minimal. Stand a safe distance from nest and not directly underneath. (Remove protective cap.) Hold can upright and point spray opening toward nest with wind at your back. Spray until nest is thoroughly saturated to kill the entire nest population. Residual action will kill wasps and hornets returning to the nest. Wait at least 24 hours before removing the nest. To kill wasps, hornets

## Aerosol Products

### Nu-Kill® II



and yellow jackets under eaves, screens, window frames, doors, patios, cracks, holes or crevices wherever insects are noticed, spray for about 1 - 2 seconds or until surfaces are moist. To prevent entrance of insects into the house, spray around points where insects enter such as outside plumbing, faucets, etc., and cracks and crevices in foundation as well as along sills and ledges. Do not apply to pets or contaminate feed, foodstuff, dishes or utensils. Avoid spraying vegetation directly.

Do not use in commercial food/feed handling establishments, restaurants or other sites where food/feed is commercially prepared, processed or served. Not for use in Federally inspected Meat and Poultry Plants. Do not use in commercial food/feed handling establishments, restaurants or other sites where food/feed is commercially prepared, processed or served. Not for use in Federally inspected Meat and Poultry Plants.

**CRAWLING INSECTS OUTDOORS:** Remove all exposed food and cooking utensils. Cover all food handling surfaces or wash thoroughly after treatment and before use. Do not use on edible crops. Point spray nozzle away from face. Hold can on slight downward angle. For best results, spray when air is calm. Spray with wind if breeze is blowing.

**PESTS ON OUTSIDE SURFACES OF BUILDINGS:** For control of building infestation due to **ANTS, CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, and SOWBUGS (PILLBUGS):** Spray foundation of building where insects

are active and may find entrance up to a height of 2 to 3 feet. Apply as a residual spray to outside surfaces of buildings where insects tend to congregate. Spray areas include (but are not limited to) screens, window frames, eaves, porches, patios, garages and refuse dumps.

**CRAWLING INSECTS - ANTS, COCKROACHES** (including ASIAN COCKROACHES), **CRICKETS, CENTIPEDES, and SILVERFISH:** Spray infested surface of patio or picnic area, hitting as many insects as possible. Also spray legs of tables and chairs to repel ants. Spray ant hills for best control.

**ANTS, LONE STAR TICKS, DOG TICKS, CRICKETS, FLEAS and EARWIGS:** Apply thoroughly to infestations in bushes, grass or weeds. For ants, spray freely around ant trails and hills.

EPA REG. NO. 10807-196-65516 EPA EST. NO. 10807-GA-1

Read and understand the product's label and Material Safety Data Sheet ("MSDS") for precautionary and first aid information.

The MSDS is available on the Nu-Calgon website at [www.nucalgon.com](http://www.nucalgon.com).

